

DRAGSTER TANK-MIX COMPATIBILITY IN FORAGE MAIZE

Must be applied with an 95.0 % w/w oil (rapeseed triglycerides) adjuvant i.e. Codacide or Vivolt

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW, especially Mixing Instructions

This version supersedes all previous editions

Normal = Physically Compatible	<i>Italics = Physically Compatible with Continuous Agitation</i>	Red on Blue = Physically Incompatible
BOLD CAPITALS = Physically & Biologically Compatible	Bold Italics = Biologically Compatible with Continuous Agitation	RED ON BLUE = CROP INJURY - INCOMPATIBLE
	Black on Orange = Do not mix as bacterial survival is severely impacted	

ADJUVANTS

CODACIDE (1.0 l/ha)	Vivolt 0.2%
---------------------	-------------

BIOSTIMULANTS

Blue N + Codacide	Blue N + Vivolt
-------------------	-----------------

HERBICIDES

Botiga	Elumis	Shield Pro + Starane Hi-Load HL + Vivolt*
CALLISTO*	Oceal + Codacide/Vivolt	Starane Hi-Load HL + Codacide*
DIVA + CODACIDE / VIVOLT	SHIELD PRO + CODACIDE	Wing-P + Codacide/Vivolt
Dual Gold + Codacide/Vivolt	Shield Pro + Starane Hi-Load HL + Codacide*	

TRACE ELEMENTS - tested in mix with Codacide (1.0 l/ha OR Vivolt 0.2%)

Amazinc	Gro-Plan P	Phosamco Bio
Biotrac	Hadron	Program
Bortrac	Intracrop Status	Prosper Plus
Brassitrel Pro	Kinsidro Grow+	Route
Canola Bio	Magflo 300	Safe N
Cearum	Magphos	Seniphos
Captrel	Maize Boost	Stopit
Croplift Pro	Mancozin	Tauron
Ferleaf	Mancufo	Thiotrac
Foliar Potash	Mantrac Pro	Zintrac
Glytrei MnP	Molytrac	
Gramitrel	Nutrino P	

COMBINATION MIXES

HERBICIDES PLUS TRACE ELEMENTS

CALLISTO + MAIZE BOOST*	Shield Pro + Starane Hi-load HL + Kinsidro Grow+ + Codacide	Starane Hi-load HL + Hadron + Kinsidro Grow+ + Vivolt
Gyo + Kinsidro Grow+ + Codacide	Shield Pro + Starane Hi-load HL + Kinsidro Grow+ + Vivolt	Starane Hi-load HL + Kinsidro Grow+ + Codacide
Gyo + Kinsidro Grow+ + Vivolt	Starane Hi-load HL + Hadron + Kinsidro Grow+ + Codacide	Starane Hi-load HL + Kinsidro Grow+ + Vivolt

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower's risk.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

NOTES:

Do not use with an adjuvant due to Callisto label restrictions. Requires the addition of an anti-foam agent.

* Do not treat if the crop is under stress.

MIXING INSTRUCTIONS:

Half fill the spray tank with clean water and start the agitation.

Pour in the required amount of DRAGSTER.

Rinse any empty containers thoroughly and add rinsings to the spray tank.

Where tank-mixes are used, and unless directed otherwise, the preferred order of addition of products to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product should be added separately to the half-full sprayer and be fully dispersed before the addition of the next product.

Add the remainder of the water.

Finally, add the required amount of adjuvant and continue agitation.

Maintain agitation until spraying is completed.

Spray solution MUST be used on the day of mixing and must NOT be left overnight.

Follow manufacturer recommendations regarding correct use of the induction bowl.

Please note:

Direct contact of WGs with non-diluted tank-mix partners must be avoided.

The above supersedes the instruction on the Codacide label and so negates the need for pre-mixing

KEY CHANGES:

Starane Hi-Load HL now approved as a mix partner; 21May 25 - addition of Hadron, Kinsidro Grow+ and herbicide mixes involving them.

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL
Corteva Agriscience™, Unit H, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG6 8HB, UK
All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
Dragster contains rimsulfuron + thifensulfuron
Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk