

**PRINCIPAL FORTE TANK-MIX COMPATIBILITY IN FORAGE MAIZE**  
**Must be applied with an 95.0 % w/w oil (rapeseed triglycerides) adjuvant i.e. Codacide**

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW  
 This version supersedes all previous editions

|   |  |  |
|---|--|--|
| <b>BOLD CAPITALS</b> = Physically & Biologically Compatible | <i>(Italics in brackets)</i> = Sprayable with Continuous Agitation                       | Red on Blue = PHYSICALLY INCOMPATIBLE    |
| Normal = Physically Compatible                              | <b>(BOLD ITALIC IN BRACKETS)</b> = Biologically compatible - maintain constant agitation | <b>RED CAPITAL ON BLUE</b> = CROP INJURY |

|                  |  |  |  |  |
|------------------|--|--|--|--|
| <b>Adjuvants</b> |  |  |  |  |
| <b>CODACIDE</b>  |  |  |  |  |

|                   |  |  |  |  |
|-------------------|--|--|--|--|
| <b>Herbicides</b> |  |  |  |  |
| <i>Botiga</i>     |  |  |  |  |
| <b>CALLISTO #</b> |  |  |  |  |

|   |                        |                        |  |  |
|---|------------------------|------------------------|--|--|
| <b>Trace Elements - tested in mix with Codacide 0.5% ONLY</b> |                        |                        |  |  |
| <i>(Amazinc)</i>  | <i>(Ferleaf)</i>       | <i>(Mantrac Pro)</i>   |  |  |
| <i>(Biotrac)</i>  | <i>(Foliar Potash)</i> | <i>(Molytrac)</i>      |  |  |
| <i>(Bortrac)</i>  | <i>(Glytrei MnP)</i>   | <i>(Phosamco Bio )</i> |  |  |
| <i>(Brassitrel Pro)</i>                                       | <i>(Gramitre)</i>      | <i>(Safe N)</i>        |  |  |
| <i>(Calfite + Luxor)</i>                                      | <i>(Magflo 300)</i>    | <i>(Senphos)</i>       |  |  |
| <i>(Canola Bio)</i>   | <i>(Magphos)</i>       | <i>(Stopt)</i>         |  |  |
| <i>(Coptrel)</i>  | <i>(Maize Boost)</i>   | <i>(Thiotrac)</i>      |  |  |
| <i>(Craplift Pro)</i>   | <i>(Mancozin)</i>      | <i>(Zintrac)</i>       |  |  |
| <i>(Equilibrium)</i>  | <i>(Mancufla)</i>      |                        |  |  |

|                                    |  |  |  |  |
|------------------------------------|--|--|--|--|
| <b>COMBINATION MIXES</b>           |  |  |  |  |
| <b>HERBICIDES + TRACE ELEMENTS</b> |  |  |  |  |
| <b>CALLISTO + MAIZE BOOST# *</b>   |  |  |  |  |

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower's risk.

**Important Note:** Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Technical Hotline on 0800 689 8899

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, PH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Spray able with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

**Notes:**

# Do not use with an adjuvant, due to Callisto label restrictions.Requires the addition of an anti-foam agent.

\* Do not apply if the crop is under stress.

**MIXING INSTRUCTIONS.**

Half fill the spray tank with clean water and start the agitation.

Pour in the required amount of PRINCIPAL FORTE.

Rinse any empty containers thoroughly and add rinsings to the spray tank.

Where tank mixes are used, and unless directed otherwise, the preferred order of addition of products to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product should be added separately to the half-full sprayer and be fully dispersed before the addition of the next product.

Add the remainder of the water.

Finally, add the required amount of adjuvant and continue agitation.

Maintain agitation until spraying is completed.

Spray solution MUST be used on the day of mixing and must NOT be left overnight.

Follow manufacturer recommendations regarding correct use of the induction bowl.

**Please note:**

Direct contact of WGs with non-diluted tank mix partners must be avoided.

The above supersedes the instruction on the Codacide label and so negates the need for pre-mixing

**KEY CHANGES**

New List; 23rd April -added Equilibrium and Calfite + Luxor

**USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.**

**FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL**

Corteva Agriscience™, CPC2 Capital Park, Fulbourn, Cambridge, Cambridgeshire, CB21 5XE. Tel: +44 (0) 1462 457272.

All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

Principal Forte contains dicamba + rimsulfuron + thifensulfuron  
 Technical Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk