

**Leystar is a selective herbicide for new sown leys, grass for seed, established grass, maize, cereals and cereals undersown with grass.**

## Key facts

Product Registration Number:	MAPP No. 21140
Active Ingredient:	144 g/l fluroxypyr + 80 g/l clopyralid + 2.5 g/l florasulam (HRAC Groups 2 & 4 Herbicides)
Pack Size:	2 litres
Formulation:	Emulsifiable concentrate (EC)
Maximum Individual Dose:	2.0 l/ha
Maximum Total Dose:	2.0 l/ha
Application Timing:	1 <sup>st</sup> February to 30 <sup>th</sup> September
Water Volumes:	200 l/ha
Spray Quality:	Medium as defined by BCPC
Buffer Zone:	5 m reducible aquatic buffer 5 m, although a LERAP can allow this to be reduced
Rainfastness:	2 hours

## Key benefits

- 3 actives delivering effective translocation and wide spectrum weed control.
- Gives excellent control of creeping and meadow buttercups, dandelions, chickweed and all species of thistles.
- Will also give good control of docks:
  - Control will be less than Doxstar® Pro but better than straight fluroxypyr when application is made in more variable temperature conditions.
- Will also give good season-long control of thistles:
  - Control will be less long-term than Thistlex®, but in optimal application conditions control will be high. However, if timing is compromised or weather not ideal and root stock significant then some limited re-growth may occur the following year.
- Handy 1 hectare pack when used at the 2.0 l/ha established grass rate.
- Good grass safety.
- Good versatility with use in new sown leys, grass grown for seed, maize, cereals and cereals undersown with grass.
- Suitable for use in grazed grassland.

## Best use advice - grass

- For best results apply to weeds which are actively growing.
- Weeds not present at application will not be controlled.
- Do not roll or harrow grass for 10 days before or 7 days after application.
- Do not spray where clover or other legumes are a valuable constituent of the sward.
- Grass seed can be stitched in 1 month after application.
- Clover can be stitched in 3 months after application.
- Grazing can resume just 7 days after application
  - This may need to be longer if foliage of poisonous weeds such as ragwort are present.
- To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, using a proprietary sprayer cleaner (i.e., All Clear Extra).

Stewardship

- Labels have now changed for clopyralid-containing products used on grassland (broadacre treatments). The changes on the labels are important and we would ask that the restrictions on the labels are observed.
- This update is to mitigate against the risk of herbicide residues ending up in manures and composts which might be used where sensitive plants are grown. More details can be found in a dedicated Topic Sheet, in our Corteva Forage App.

Weed spectrum

Control of Annual Weeds with Leystar 1.0 l/ha					
Bindweed (black)	◆ S up to 4TL	Fool's parsley	R	Pale persicaria	◆ MS up to 2TL
Bristly ox-tongue	◆ S	Forget-me-not	◆ S up to 4TL	Poppy	◆ MS up to 4TL
<b>Charlock</b>	<b>S up to 1TL</b>	Fumitory	◆ MS up to 2TL	Redshank	◆ MS up to 2TL
Chickweed (common)	◆ S up to 100 mm and BF	Groundsel	◆ MS up to 2TL	Scarlet pimpernel	R
Chickweed (mouse-eared)	◆ MS up to 100mm and BF	Hemp-nettle	◆ S up to 4TL	<b>Shepherd's purse</b>	<b>S up to 4TL</b>
<b>Cleavers</b>	<b>S up to 200 mm and BF</b>	Himalayan balsam		Speedwells	R
Corn chamomile	◆ S up to 150 mm and BF	Knotgrass	◆ S up to 4TL	Spurrey	◆ MS up to 2TL
Corn marigold	◆ S up to 6TL	<b>Mayweeds</b>	<b>S up to 1TL</b>	Vol OSR	S up to 4TL
Cranesbill	R	Nettle (small)	R	Wild radish (runch)	S up to 4TL and 50mm
Dead nettles	◆ MS up to 2TL	Nightshade (black)	◆ MS	Yellow rattle	
Fat-hen	R	Orache	R		
Control of Perennial Weeds with Leystar 2.0 l/ha					
Bindweed (field)	◆ S	Gorse	R	Old man's beard	
Bracken		Ground elder	R	<b>Plantain (greater)</b>	<b>MS up to Rosette and BF</b>
Bramble	R	Ground ivy		<b>Plantain (ribwort)</b>	<b>MS up to Rosette and BF</b>
Broom	R	Hawthorn	R	Ragwort	R
Burdock		Hemlock	◆ MR	Rosebay willowherb	
<b>Buttercups</b>	<b>S up to BF</b>	Hogweed	R	Rushes	R
Cinquefoil (creeping)	◆ MR	Horsetail (Equisetum)	R	Self-heal	
Clover, trefoil	◆ S	Japanese knotweed	R	Silverweed	
Coltsfoot	◆ MS	Knapweed (common)	◆ MR	Sorrel (common)	◆ MS up to Rosette and BF
Cow parsley	R	Lesser celandine		Thistles	◆ S up to 150 mm and BF
<b>Daisy (common)</b>	<b>S up to Rosette and BF</b>	Mallow (common)	◆ MS	Vetch, tare	◆ S
Daisy (ox-eye)	◆ MR	Medick (black)		Yarrow	◆ S
<b>Dandelion</b>	<b>S up to Rosette and BF</b>	Mugwort (common)		Yellow/Flag Iris	
<b>Docks</b>	<b>MS up to 200 mm and BF</b>	Nettle (common)	R		

**Key:** Label weeds in **bold**. TL – True leaves, B4F – Before flowering, \* - Season-long control, ◆ - indicates information based on anecdotal or limited data and is only indicative and should not be considered as a recommendation for use on the part of Corteva Agriscience. The user assumes full responsibility for use on these weeds.

	Susceptible		Moderately susceptible
	Moderately resistant		Resistant
	No Information		