

WATER VOLUME AND APPLICATION

Apply ZORVEC ENDAVIA in at least 200 L water/hectare using a MEDIUM quality spray (as defined by BCPC) to give good even coverage of the crop. Higher water volumes should be used in dense crops to improve coverage.

COMPATIBILITY

Products should only be tank-mixed if each product can be applied within the label recommendations for its use. All other requirements and restrictions on the partner product label(s) should be adhered to. Consult your Corteva distributor for further details

SPRAY TANK CLEANING

Immediately after application, completely drain spray equipment. Thoroughly rinse sprayer and flush the hoses, boom nozzles with clean water.

Loosen and physically remove visible deposits.

Remove and clean nozzles, screens and strainers.

Flush the entire system with clean water.

Take all necessary safety precautions when cleaning equipment.

Do not clean near wells or water sources.

Consult label tank cleanup procedures for all tank-mix partners and be sure to use the most rigorous procedure recommended

Dispose of waste rinse water in accordance with local regulations.

NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any

damage or injury whatsoever arising from their storage, handling, application or use. Since the occurrence of resistance cannot be forecast, neither Du Pont (UK) Limited nor its distributors can accept responsibility for any loss or damage to crops caused by the failure of ZORVEC ENDAVIA to control resistant strains.

All trademarks duly acknowledged.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

Operators must wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

Environmental protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place

KEEP OUT OF REACH OF CHILDREN

EMPTY CONTAINER COMPLETELY and dispose of safely

DO NOT RE-USE CONTAINER for any purpose.

Batch No.

Manufactured in the EU

FOR PROFESSIONAL USE ONLY

SHAKE WELL BEFORE USE

To access the Safety Data Sheet for this product scan the code, visit www.corteva.co.uk or alternatively contact your supplier



Zorvec Endavia®

FUNGICIDE

MAPP 19407

An oil dispersion fungicide containing 70 g/L benthiavalicarb and 30 g/L oxathiapiprolin for disease control in potatoes, onions, garlic and shallots.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Authorisation holder:

Du Pont (U.K.) Limited, Crop Protection Products Kings Court,

London Road, Stevenage Hertfordshire., SG1 2NG

Email: ukhotline@corteva.com – Tel 0800 689 8899

24 HOUR EMERGENCY TELEPHONE NUMBER : 0161 884 1235

National Poison Centre: 111 (England and Wales) 08454 24 24 24 (Scotland)

©™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

CONTENTS : 5L e

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crop	Max. Individual Dose	Max. No. of applications	Latest time of Application
Potato	0.4 L/ha	4 per crop	7 days pre-harvest*
Onion, Garlic, Shallot	0.5 L/ha	3 per crop	28 days pre-harvest

* See Directions for Use

Other Specific Restrictions

- On potatoes, a minimum interval of 7 days must elapse between application and initial haulm destruction and/or harvest, whichever comes first.
- The minimum interval between applications is 7 days

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

K-47019/2002 - UNITED KINGDOM

Lot N° : see on the pack



3 576960 069083

Safety Information

ZORVEC ENDAVIA

Contains 70 g/L benthiavalicarb and 30 g/L oxathiapiprolin

May cause an allergic skin reaction.

Suspected of causing cancer.

Toxic to aquatic life with long lasting effects

Avoid breathing spray.

Wear protective gloves/protective clothing/eye protection/face protection

Do not eat, drink or smoke when using this product.

IF ON SKIN: Wash with plenty of soap and water.

IF ON EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

If skin irritation or a rash occurs: Get medical advice/attention

Wash contaminated clothing before reuse.

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

Collect spillage.


Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

To avoid risks to human health and the environment, comply with the instructions for use

WARNING



OPEN HERE

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

- Avoid application to any crop suffering from stress as a result of drought, waterlogging, low temperatures, nutrient or lime deficiency or other factors reducing crop growth.

DISEASE CONTROL

ZORVEC ENDAVIA combines two active ingredients: oxathiapiprolin and bentiavalicarb with different modes of action. Oxathiapiprolin is listed by FRAC Fungicide Resistance Action Committee (FRAC) as Code 49. Bentiavalicarb is a Carboxylic Acid Amides (CAA) fungicide, classified under FRAC Code 40

ZORVEC ENDAVIA should be used as part of a season long programme of correctly timed sprays commencing before disease infection. Application should be made to dry foliage.

Potatoes

ZORVEC ENDAVIA can be used on all varieties of early and maincrop potatoes. A season long disease control programme is required to limit late blight development. This should include a haulm dessicant. Crops should not be harvested until at least 7 days after all green tissue has died.

Apply preventatively before the first visible signs of blight in the crop. Commence treatment with ZORVEC ENDAVIA at the first blight warning or when local weather conditions are favourable for disease development. Applications should be repeated at 7-10 day intervals depending on disease pressure. As disease pressure increases, the interval should be shortened.

Apply ZORVEC ENDAVIA at 0.4 L/ha. No more than 3 applications of ZORVEC ENDAVIA may be applied consecutively

Onions, Garlic and Shallot

ZORVEC ENDAVIA can be used on all varieties from the 3-leaf stage of growth. Apply preventatively before the first visible signs of downy mildew.

Apply ZORVEC ENDAVIA at 0.5 L/ha. No more than 2 applications may be applied consecutively.

RESISTANCE

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. A pathogen is considered resistant to a fungicide if it survives a correctly applied treatment at the recommended dose and timing under normal weather conditions. Plant disease biotypes resistant to oxathiapiprolin may exist. Development of resistance can be avoided or delayed by alternating or mixing products with different modes of action.

The maximum number of recommended applications of ZORVEC ENDAVIA should not be exceeded and it should be incorporated into spray programmes that include fungicides with differing modes of action. No more than three consecutive applications of a CAA fungicide should be made. Please refer to Fungicide Resistance Action Committee (FRAC) guidelines for further details and always ensure good resistance management practice is followed to reduce the incidence of resistance developing.

MIXING

Spraying equipment should be thoroughly cleaned before spraying. ZORVEC ENDAVIA mixes easily with water and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is complete.